

SUPERSYMMETRY AND BLACK HOLES *

RENATA KALLOSH

*Physics Department, Stanford University,
Stanford, CA 94305*

We consider static black holes, which are bosonic solutions of supersymmetric theories. We will show that supersymmetry provides a natural framework for a discussion of various properties of such static black holes. The most fundamental property of simple global supersymmetry, non-negativeness of the energy, is generalized in the black hole theory to cosmic censorship. The SUSY classification of static black holes will be presented in terms of central charges of supersymmetry algebra. The mass, the temperature and the entropy of these black holes are simple functions of supersymmetry charges. The extreme black holes have zero temperature and some unbroken supersymmetries.

Simple $N = 1$ global supersymmetry algebra relates supercharges to the Hamiltonian $H = P^0$.

$$\{Q_\alpha, \bar{Q}_{\dot{\beta}}\} = (\sigma_\mu)_{\alpha\dot{\beta}} P^\mu . \quad (1)$$

This implies that the spectrum of the Hamiltonian is semipositive definite since $\bar{Q}$ is Hermitean adjoint of Q and we have

$$H = P^0 = (\bar{Q}_1 Q_1 + Q_1 \bar{Q}_1 + \bar{Q}_2 Q_2 + Q_2 \bar{Q}_2) \equiv |Q|^2 \geq 0 . \quad (2)$$

Local supersymmetry does not require positivity of energy, in general. However, if the curved space is asymptotically flat, states at infinity may be described by the multiplets of global supersymmetry. In particular, a generalization of the positiveness of the energy has been found for static black holes, considered as bosonic solutions of supersymmetric theories in ^{1,2}. Axion-dilaton $U(1)^2$ black holes of a mass M in four-dimensional theory are associated, as shown in ref. ² with extended $N = 4$ supersymmetry algebra with central charges.

$$\{Q_\alpha^I, Q_{\dot{\beta}}^{*J}\} = (\sigma_\mu)_{\alpha\dot{\beta}} P^\mu \delta^{IJ} , \quad (3)$$

$$\{Q_\alpha^I, Q_\beta^J\} = \epsilon_{\alpha\beta} Z^{IJ} , \quad Z^{IJ} = -V^{IJ} + iU^{IJ} \quad (4)$$

$$\{Q_{\dot{\alpha}}^{*I}, Q_{\dot{\beta}}^J\} = \epsilon_{\dot{\alpha}\dot{\beta}} \bar{Z}^{IJ} . \quad (5)$$

In the rest frame in a special basis this algebra describes a system of states, characterized by the mass and two complex central charges z_1 and z_2 .

$$\{q_{(1)}, q_{(1)}^*\} = 2|q_{(1)}|^2 = M - |z_1| \geq 0 ,$$

* Invited talk given at SUSY 93, Boston, April 1993.

$$\begin{aligned}
\{q_{(2)}, q_{(2)}^*\} &= 2|q_{(2)}|^2 = M - |z_2| \geq 0, \\
\{q_{(3)}, q_{(3)}^*\} &= 2|q_{(3)}|^2 = M + |z_1| \geq 0, \\
\{q_{(4)}, q_{(4)}^*\} &= 2|q_{(4)}|^2 = M + |z_2| \geq 0.
\end{aligned}
\tag{6}$$

(7)

The central charges are functions of electric Q and magnetic P charges of the vector field $F_{\mu\nu}^1$ and axial-vector field $F_{\mu\nu}^2$.

$$z_1 = \sqrt{2}(\Gamma_1 + i\Gamma_2), \tag{8}$$

$$z_2 = \sqrt{2}(\Gamma_1 - i\Gamma_2), \tag{9}$$

where $\Gamma = \frac{1}{2}(Q + iP)$. The energy (the mass) of the charged black hole is not only non-negative but according to supersymmetry algebra (7) it has to be greater or equal to central charges.

$$M \geq |z_1|, \quad M \geq |z_2|. \tag{10}$$

It is quite surprising that for all known to us static black holes in supersymmetric theories the limits (10), required by supersymmetry, coincide exactly with the limits on the parameters of the black holes required by the cosmic censorship conjecture. The necessary condition is the possibility of embedding the bosonic action into a supersymmetric action. For example, for the black holes which are obtained from the action where dilaton interacts with vector fields as $e^{2a\phi}F_{\mu\nu}^2$ and a is an arbitrary parameter, we do not have any information of that kind except when $a = 1$ (or $\sqrt{3}$) and supersymmetric embedding is known.

The most important parameters describing the black holes are its mass and charges. In addition, one usually calculates the Hawking temperature and the Bekenstein-Hawking entropy of a black hole, which are usually some functions of the mass and charges. We have found that all these parameters of a black hole can be expressed in terms of the supercharges, describing this system. In particular, the mass is given by one half of the sum over all four supercharges $|q_{(1)}|, |q_{(2)}|, |q_{(3)}|, |q_{(4)}|$.

$$M = \frac{1}{2} \sum_{i=1}^{i=4} |q_{(i)}|^2. \tag{11}$$

The product of the temperature and entropy is proportional to the product of all four supercharges.

$$S T = \frac{4 \prod_{i=1}^{i=4} |q_{(i)}|}{\sum_{j=1}^{j=4} |q_{(j)}|^2} \tag{12}$$

The temperature is also a function of supercharges.

$$T = \frac{1}{\pi \sum_{j=1}^{j=4} |q_{(j)}|^2} \frac{\prod_{i=1}^{i=4} |q_{(i)}|}{(|q_{(1)}q_{(3)}| + |q_{(2)}q_{(4)}|)^2}. \tag{13}$$

The Hawking temperature of the black hole, according to this equation, vanishes if one of the eigenvalues of the 4 supercharges is zero. However, if two supercharges approach zero eigenvalues, the limiting temperature depends on the order in which this limit is taken. If first we assume that for example $|q_{(1)}| \rightarrow |q_{(2)}|$ and $|q_{(1)}| \rightarrow |q_{(2)}|$, the limiting temperature equals $T = \frac{1}{8\pi M}$. This correspond to

the limit from $U(1)^2$ black hole with two vector fields to $U(1)$ black hole with one vector field. If however one first take the limit $|q_{(1)}| \rightarrow 0$ at fixed $|q_{(2)}|$, the temperature goes to zero and remains zero when the limit $|q_{(2)}| \rightarrow 0$ is taken afterwards.

The entropy can be represented as a function of supercharges:

$$S = 4\pi(|q_{(1)}q_{(3)}| + |q_{(2)}q_{(4)}|)^2. \quad (14)$$

As it follows from this equation, the entropy vanishes when $|q_{(1)}| = |q_{(2)}| = 0$. The zero eigenvalue of the supercharge indicates the existence of unbroken supersymmetries.

The fact that the mass, the temperature and the entropy of static black holes can be completely described in terms of the eigenvalues of the supercharges is quite remarkable! This is a property of all static black holes and not just of extreme black holes which have some unbroken supersymmetries.

Supersymmetry provides a simple classification of four-dimensional static black holes, starting with classical Schwarzschild and Reissner-Nordström black holes up to axion-dilaton black holes.

Our conventions are those of ². We will use the complex scalar $\lambda = a + ie^{-2\phi}$, where a is the axion field and ϕ is the dilaton field. We also have two $U(1)$ vector fields $A_\mu^{(n)}$, $n = 1, 2$.

We find it convenient to define the $SL(2, R)$ -duals[†] to the fields $F_{\mu\nu}^{(n)} = \partial_\mu A_\nu^{(n)} - \partial_\nu A_\mu^{(n)}$,

$$\tilde{F}^{(n)} = e^{-2\phi} \star F^{(n)} - ia F^{(n)}, \quad (15)$$

in terms of which the bosonic part of $N = 4$ supergravity action reads

$$S = \frac{1}{16\pi} \int d^4x \sqrt{-g} \left\{ -R + \frac{1}{2} \frac{\partial_\mu \lambda \partial^\mu \bar{\lambda}}{(\text{Im } \lambda)^2} - \sum_{n=1}^2 F_{\mu\nu}^{(n)} \star \tilde{F}^{(n)\mu\nu} \right\}. \quad (16)$$

In terms of the component fields, we have

$$S = \frac{1}{16\pi} \int d^4x \sqrt{-g} \left\{ -R + 2(\partial\phi)^2 + \frac{1}{2} e^{4\phi} (\partial a)^2 - e^{-2\phi} \sum_{n=1}^2 (F^{(n)})^2 + ia \sum_{n=1}^2 F^{(n)} \star F^{(n)} \right\}. \quad (17)$$

The advantage of using $\tilde{F}^{(n)}$ is that the equations of motion imply the local existence of N real vector potentials $\tilde{A}^{(n)}$ such that

$$\tilde{F}^{(n)} = i d\tilde{A}^{(n)}. \quad (18)$$

The analogous equation $F^{(n)} = dA^{(n)}$ is not a consequence of equations of motion but a consequence of the Bianchi identity. If the time-like components $A_t^{(n)}$ play the role of electrostatic potentials, then the $\tilde{A}_t^{(n)}$ will play the role of magnetostatic potentials. The $SL(2, R)$ duality transformations consist in the mixing of $A^{(n)}$ with

[†]The spacetime duals are $\star F^{(n)\mu\nu} = \frac{1}{2\sqrt{-g}} \epsilon^{\mu\nu\rho\sigma} F_{\rho\sigma}$, with $\epsilon^{0123} = \epsilon_{0123} = +i$.

$\tilde{A}^{(n)}$ and of equations of motion with Bianchi identities, as in the Einstein-Maxwell case.

Here we present two different kinds of static solutions to the equations of motion of the action (16), (17): spherical black-hole solutions and multi-black-hole solutions, both with nontrivial axion, dilaton and $U(1)$ fields. All the previously known solutions of these kinds (Schwarzschild, (multi-) Reissner-Nordström, the electric and magnetic dilaton black holes, the electric-magnetic axion-dilaton black holes of refs.^{3,2} are particular cases of them.

The static spherically symmetric black-hole solutions are²

$$\begin{aligned} ds^2 &= e^{2U} dt^2 - e^{-2U} dr^2 - R^2 d\Omega^2, \\ \lambda(r) &= \frac{\lambda_0 r + \bar{\lambda}_0 \Upsilon}{r + \Upsilon}, \\ A_t^{(n)}(r) &= e^{\phi_0} R^{-2} [\Gamma^{(n)}(r + \Upsilon) + c.c], \\ \tilde{A}_t^{(n)}(r) &= -e^{\phi_0} R^{-2} [\Gamma^{(n)}(\lambda_0 r + \bar{\lambda}_0 \Upsilon) + c.c], \end{aligned} \quad (19)$$

where

$$\begin{aligned} e^{2U}(r) &= R^{-2}(r - r_+)(r - r_-), \quad r_{\pm} = M \pm r_0, \\ R^2(r) &= r^2 - |\Upsilon|^2, \quad r_0^2 = M^2 + |\Upsilon|^2 - 4 \sum_{n=1}^N |\Gamma^{(n)}|^2. \end{aligned} \quad (20)$$

We define the parameters of our solutions in terms of the asymptotic behavior ($r \rightarrow \infty$) of different complex fields

$$\begin{aligned} g_{tt} &\sim 1 - \frac{2M}{r}, & \lambda &\sim \lambda_0 - ie^{-2\phi_0} \frac{2\Upsilon}{r}, \\ F_{tr} &\sim \frac{e^{+\phi_0} Q}{r^2}, & {}^*F_{tr} &\sim \frac{ie^{+\phi_0} P}{r^2}. \end{aligned} \quad (21)$$

The real axion (Δ), dilaton (Σ), electric (Q) and magnetic (P) charges, and the asymptotic values of the axion (a_0) and dilaton (ϕ_0) are

$$\Upsilon = \Sigma - i\Delta, \quad \Gamma = \frac{1}{2}(Q + iP), \quad \lambda_0 = a_0 + ie^{-2\phi_0}. \quad (22)$$

In every black hole described by our solutions the charge of the complex scalar Υ is related to the electromagnetic charges by

$$\Upsilon = -\frac{2}{M} (\bar{\Gamma}_n)^2. \quad (23)$$

The singularity is hidden under the horizon if $r_0^2 > 0$, and it is hidden or coincides with it (but still is invisible for an external observer) if $r_0 = 0$. The conditions $r_0^2 \geq 0$ and $M \geq |\Upsilon|$ can be related to supersymmetry bounds². All solutions given above have the entropy

$$S = \pi(r_+^2 - |\Upsilon|^2). \quad (24)$$

When all supersymmetric bounds are saturated, i.e. $r_+ = M = |\Upsilon|$, the objects described by this solution have zero area of the horizon and vanishing entropy. In this sense, such black holes (*holons*) behave as elementary particles.

The electric, magnetic, dilaton and axion charges can be expressed in terms of the central charges z_1, z_2 and the mass M :

$$\Gamma_1 = \frac{1}{2\sqrt{2}}(z_1 + z_2) , \quad (25)$$

$$\Gamma_2 = \frac{1}{2\sqrt{2}i}(z_1 - z_2) , \quad (26)$$

$$\tilde{\Upsilon} = \Sigma + i\Delta = -\frac{z_1 z_2}{M} . \quad (27)$$

$$(28)$$

Thus our axion-dilaton black holes have the following “hair”: the mass M , two complex central charges z_1, z_2 (through which electric, magnetic, dilaton and axion charges are expressed) and asymptotic value at infinity of the dilaton and of the axion field $\lambda_0 = a_0 + e^{-2\phi_0}$.

The temperature of a whole family of axion-dilaton black holes is

$$T = \frac{1}{2\pi M} \frac{\sqrt{(M^2 - |z_1|^2)(M^2 - |z_2|^2)}}{\left[\sqrt{M^2 - |z_1|^2} + \sqrt{M^2 - |z_2|^2}\right]^2} . \quad (29)$$

The entropy, which equals one quarter of the area of the horizon, is

$$S = \pi \left[\sqrt{M^2 - |z_1|^2} + \sqrt{M^2 - |z_2|^2}\right]^2 . \quad (30)$$

The parameter r_0 , which vanishes when the black hole becomes extremal, is given by the product of supersymmetry bounds:

$$r_0^2 = \frac{(M^2 - |z_1|^2)(M^2 - |z_2|^2)}{M^2} . \quad (31)$$

SUSY CLASSIFICATION OF STATIC BLACK HOLES will be given in terms of central charges and asymptotic values of dilaton-axion field at infinity.

Schwarzschild solution corresponds to $z_1 = z_2 = \lambda_0 = i$. Thus we get $r_0 = M$, $r_+ = 2M$, $r_- = 0$, the vector and dilaton-axion fields are absent in this solution. The temperature and entropy are obtained from eqs. (29), (30).

$$T_{Sch} = \frac{1}{8\pi M} ,$$

$$S_{Sch} = 4\pi M^2 . \quad (32)$$

Reissner-Nordström solution is given by $z_1 = \lambda_0 = i$, $z_2 = q$. We have $r_0 = \sqrt{M^2 - q^2}$, $r_+ = 2M$, $r_- = 0$, the vector and dilaton-axion fields are absent in this solution. Substitution of these values of central charges into eqs. (29), (30) gives

$$T_{RN} = \frac{1}{2\pi} \frac{\sqrt{M^2 - q^2}}{\left(M + \sqrt{M^2 - q^2}\right)^2} ,$$

$$S_{RN} = \pi \left(M + \sqrt{M^2 - q^2}\right)^2 . \quad (33)$$

Extreme Reissner-Nordström black holes with $M = q$ have zero temperature and non-zero entropy, $N = 1$ supersymmetry being unbroken¹.

$U(1)$ **Charged axion-dilaton black holes**³ have $|z_1| = |z_2| = |Q/\sqrt{2}|$. The corresponding expressions for the temperature and the entropy are

$$\begin{aligned} T_{U(1)} &= \frac{1}{2\pi M} \frac{M^2 - |Q/\sqrt{2}|^2}{4(M^2 - |Q/\sqrt{2}|^2)} = \frac{1}{8\pi M} , \\ S_{U(1)} &= 4\pi \left(M^2 - |Q/\sqrt{2}|^2 \right)^2 . \end{aligned} \quad (34)$$

In the extreme limit $M = |Q/\sqrt{2}|$ $N = 2$ supersymmetry is restored. The temperature remains equal to the temperature of the Schwarzschild black hole (however, see the discussion above where it is explained that this temperature takes different values if the limit to extreme is taken starting from $U(1) \times U(1)$ configuration) and the entropy vanishes.

Finally consider an $U(1) \times U(1)$ **four-dimensional axion-dilaton black holes**, with one vector and one axial vector field. The central charges in terms of electromagnetic charges are given by

$$\begin{aligned} |z_1|^2 &= \frac{1}{2}(P_1^2 + Q_1^2 + P_2^2 + Q_2^2) - E , \\ |z_2|^2 &= \frac{1}{2}(P_1^2 + Q_1^2 + P_2^2 + Q_2^2) + E , \end{aligned} \quad (35)$$

where

$$E \equiv Q_1 P_2 - Q_2 P_1 . \quad (36)$$

Q_1, P_1 and Q_2, P_2 are electric and magnetic charges of a vector and axial-vector fields, respectively. The entropy and temperature of these solutions are given in eqs. (29), (30).

Extreme charged $U(1) \times U(1)$ axion-dilaton black holes have a restored $N = 1$ supersymmetry. Extreme solution $M = |z_1|$ or $M = |z_1|, |z_1| \neq |z_2|$ has zero temperature and non-zero entropy,

$$\begin{aligned} T &= 0 , \\ S &= \pi (|z_1|^2 - |z_2|^2) = \pi(Q_1 P_2 - Q_2 P_1) . \end{aligned} \quad (37)$$

There are different ways to reach the restoration of $N = 2$ supersymmetry. One possibility is to consider the next step $|z_1| \rightarrow |z_2|$. In this particular limit

$$\begin{aligned} T &= 0 , \\ S &= 0 . \end{aligned} \quad (38)$$

and the system seems to approach the ground state with zero temperature and entropy.

The analogous situation was discussed at length and illustrated in figures in ref. 2, where we plotted both temperature and entropy either as the functions of central charges at fixed mass or as the function of mass at a given central charges. All figures presented there for the dilaton solution apply to the more general axion-dilaton solutions under the condition that instead of the central charges one should

use the moduli of complex central charges, since only in absence of axion are the central charges real.

Our second kind of solutions describe axion-dilaton extreme multi-black-hole solutions². **Multi-black-holes have unbroken supersymmetries.** The fields are

$$\begin{aligned}
ds^2 &= e^{2U} dt^2 - e^{-2U} d\vec{x}^2, & e^{-2U}(\vec{x}) &= 2 \operatorname{Im}(\mathcal{H}_1(\vec{x}) \overline{\mathcal{H}_2(\vec{x})}), \\
\lambda(\vec{x}) &= \frac{\mathcal{H}_1(\vec{x})}{\mathcal{H}_2(\vec{x})}, \\
A_i^{(n)}(\vec{x}) &= e^{2U}(k^{(n)}\mathcal{H}_2(\vec{x}) + c.c.), \\
\tilde{A}_i^{(n)}(\vec{x}) &= -e^{2U}(k^{(n)}\mathcal{H}_1(\vec{x}) + c.c.),
\end{aligned} \tag{39}$$

where $\mathcal{H}_1(\vec{x}), \mathcal{H}_2(\vec{x})$ are two complex harmonic functions

$$\begin{aligned}
\mathcal{H}_1(\vec{x}) &= \frac{e^{\phi_0}}{\sqrt{2}} \left\{ \lambda_0 + \sum_{i=1}^I \frac{\lambda_0 M_i + \bar{\lambda}_0 \Upsilon_i}{|\vec{x} - \vec{x}_i|} \right\}, \\
\mathcal{H}_2(\vec{x}) &= \frac{e^{\phi_0}}{\sqrt{2}} \left\{ 1 + \sum_{i=1}^I \frac{M_i + \Upsilon_i}{|\vec{x} - \vec{x}_i|} \right\}.
\end{aligned} \tag{40}$$

The horizon of the i -th (extreme) black hole is at $\vec{x}_i$ (in these isotropic coordinates the horizons look like single points), and has mass M_i , electromagnetic charge Γ_i , etc. , as can be seen by using the definitions in the Appendix in the limit $|\vec{x} - \vec{x}_i| \rightarrow \infty$. Charges without label are total charges. The constants $k^{(n)}$ are

$$k^{(n)} = -\sqrt{2} \left(\frac{\Gamma^{(n)} M + \overline{\Gamma^{(n)}} \Upsilon}{M^2 - |\Upsilon|^2} \right). \tag{41}$$

The consistency of the solution requires for every i

$$k_i^{(n)} = k^{(n)}, \quad \operatorname{Arg}(\Upsilon_i) = \operatorname{Arg}(\Upsilon). \tag{42}$$

Finally, for each i and also for the total charges, the supersymmetric Bogomolny bound is saturated:

$$M^2 + |\Upsilon|^2 - 4|\Gamma^{(n)}|^2 = 0. \tag{43}$$

For a single $U(1)$ vector field the extreme solution simplifies to ($k \equiv k^1$, $\Gamma \equiv \Gamma^1$)

$$M^2 = |\Upsilon|^2, \quad k = -\frac{\sqrt{2} \Gamma}{M}. \tag{44}$$

As a consequence of all the identities obeyed by the charges, it is possible to derive the following expression of equilibrium of forces between two extreme black holes²:

$$M_1 M_2 + \Sigma_1 \Sigma_2 + \Delta_1 \Delta_2 = Q_1 Q_2 + P_1 P_2. \tag{45}$$

If supersymmetry will be discovered in the coming experiments, it will be much more natural to study non-perturbative aspects of quantum gravity in the

context of supersymmetric theories. At the present time supersymmetric approach to quantum gravity is motivated mainly by its internal consistency and existence of natural structures describing such interesting objects as black holes. An important advantage of this approach to the theory of black holes is the possibility to obtain results which in certain cases do not acquire any quantum corrections².

In this paper we have discussed many properties of extreme and non-extreme black holes in terms of supercharges. Our main conclusion is that in supersymmetric theories evaporation of black holes is related to restoration of supersymmetry.

This work was supported by the NSF grant PHY-8612280 and in part by Stanford University.

References

1. G.W. Gibbons and C.M. Hull, *Phys. Lett.* **109B**, (1982) 190.
2. R. Kallosh, A. Linde, T. Ortín, A. Peet and A. Van Proeyen, *Phys. Rev.* **D46**, 5278 (1992);
R. Kallosh and T. Ortín, *Phys. Rev.* **D48**, 713 (1993);
R. Kallosh and T. Ortín, *Killing Spinor Identities*, preprint SU-ITP-93-16 (hep-th/9306085).
3. G. W. Gibbons, *Nucl. Phys.* **B207** (1982) 337;
G. W. Gibbons and K. Maeda, *Nucl. Phys.* **B298** (1988) 741;
D. Garfinkle, G. Horowitz and A. Strominger, *Phys. Rev.* **D43** (1991) 3140;
A. Shapere, S. Trivedi and F. Wilczek, *Mod. Phys. Lett.* **A6** (1991) 2677.